


B. Claims

The following is a complete listing of the claims, and replaces all earlier versions and listings.


1-20. (Cancelled)

21. (Currently Amended) A polyhydroxyalkanoate comprising a monomer unit of 3-hydroxy- ω -(phenylmethyl)oxy]alkanoic acid expressed by chemical formula (1):




wherein x can be one or more integers within the range shown in the chemical formula (1).

22. (Currently Amended) The polyhydroxyalkanoate according to claim 21, comprising at least one unit expressed by a chemical formula selected from the group consisting of chemical formulas (2) and (3):




wherein y and z can be one or more integers within the range shown in the chemical formulas (2) and (3), while being independent from the monomer unit expressed by the chemical formula (1).

23. (Currently Amended) The polyhydroxyalkanoate according to claim 21, comprising simultaneously, in at least a molecule thereof, the monomer of 3-hydroxy- ω -(phenylmethyl)oxy]alkanoic acid expressed by the chemical formula (1) and a unit expressed by chemical formula (4):




wherein m can be one or more integers within the range shown in the chemical formula (4), and R comprises a residue having either a phenyl structure or a

thienyl structure, or a 3-hydroxy- ω -cyclohexylalkanoic acid unit expressed by chemical formula (5):




wherein R₁ is H, CN, NO₂, halogen, CH₃, C₂H₅, C₃H₇, CF₃, C₂F₅, and C₃F₇, and k can be one or more integers within the range shown in the chemical formula (5),


wherein R in chemical formula (4), ~~i.e. a residue having either a phenyl structure or a thienyl structure,~~ is at least one group selected from the group consisting of residues




wherein R₂ is H, halogen, CN, NO₂, CH₃, C₂H₅, C₃H₇, CH=CH₂, CF₃, C₂F₅, C₃F₇, and COOR₃, ~~(wherein where R₃ represents any one selected from the group consisting of H, Na, and K), CF₃, C₂F₅, and C₃F₇,~~ and in a case where there exist a plurality of units, R₂ may be different for each unit;




wherein R₄ is selected from the group consisting of H, halogen, CN, NO₂, CH₃, C₂H₅, C₃H₇, SCH₃, CF₃, C₂F₅, and C₃F₇, and in a case where there exist a plurality of units, R₄ may be different for each unit;






wherein R₅ is selected from the group consisting of H, halogen, CN, NO₂, CH₃, C₂H₅, C₃H₇, CF₃, C₂F₅, and C₃F₇, and in a case where there exist a plurality of units, R₅ may be different for each unit;




wherein R₆ is selected from the group consisting of H, halogen, CN, NO₂, CH₃, C₂H₅, C₃H₇, (CH₃)₂-CH, (CH₃)₃-C, COOR₇, and SO₂R₈, ~~(wherein where~~ R₇ represents any one selected from the group consisting of H, Na, K, CH₃, and C₂H₅, and R₈ represents any one selected from the group consisting of OH, ONa, OK, halogen, OCH₃, and OC₂H₅); ~~CH₃, C₂H₅, C₃H₇, (CH₃)₂-CH, and (CH₃)₃-C,~~ and in a case where there exist a plurality of units, R₆ may be different for each unit;




wherein R₉ represents a substituent group on the aromatic ring, R₉ is selected from the group consisting of H, halogen, CN, NO₂, CH₃, C₂H₅, C₃H₇, (CH₃)₂-CH, (CH₃)₃-C, COOR₁₀, and SO₂R₁₁, ~~wherein~~ where R₁₀ represents any one selected from the group consisting of H, Na, K, CH₃, and C₂H₅, and R₁₁ represents any one selected from the group consisting of OH, ONa, OK, halogen, OCH₃, and OC₂H₅), ~~CH₃, C₂H₅, C₃H₇, (CH₃)₂-CH and (CH₃)₃-C~~, and in a case where there exist a plurality of units, R₉ may be different for each unit;






wherein R₁₂ is selected from the group consisting of H, halogen, CN, NO₂, CH₃, C₂H₅, C₃H₇, (CH₃)₂-CH, (CH₃)₃-C, COOR₁₃, and SO₂R₁₄, ~~wherein where R₁₃~~ represents any one selected from the group consisting of H, Na, K, CH₃, and C₂H₅, and R₁₄ represents any one selected from the group consisting of OH, ONa, OK, halogen, OCH₃, and OC₂H₅), ~~CH₃, C₂H₅, C₃H₇, (CH₃)₂-CH and (CH₃)₃-C~~, and in a case where there exist a plurality of units, R₁₂ may be different for each unit; and




wherein R₁₅ is selected from the group consisting of H, halogen, CN, NO₂, CH₃, C₂H₅, C₃H₇, (CH₃)₂-CH, (CH₃)₃-C, COOR₁₆, and SO₂R₁₇, ~~wherein where R₁₆~~ represents any one selected from the group consisting of H, Na, K, CH₃, and C₂H₅, and R₁₇ represents any one selected from the group consisting of OH, ONa, OK, halogen, OCH₃, and OC₂H₅), ~~CH₃, C₂H₅, C₃H₇, (CH₃)₂-CH and (CH₃)₃-C~~, and in a case where there exist a plurality of units, R₁₅ may be different for each unit.

24. (Previously Presented) The polyhydroxyalkanoate according to claim 21, wherein a number average molecular weight is within the range between 1000 and 1000000.

25. (Currently Amended) A method for producing a polyhydroxyalkanoate comprising, in a molecule thereof, a ~~monomer unit of 3-hydroxy- ω -(phenylmethyl)oxy]alkanoic acid~~ monomer unit expressed by chemical formula (1):




wherein x can be one or more integers within the range shown in the chemical formula (1), ~~which comprises the method comprising~~ allowing a microorganism ~~with an ability to produce a polyhydroxyalkanoate comprising in a molecule thereof the monomer unit of 3-hydroxy- ω -(phenylmethyl)oxy]alkanoic acid expressed by chemical formula (1)~~ of one or more strains selected from the group consisting of *Pseudomonas cichorii* YN2 (FERM BP-7375), *Pseudomonas cichorii* H45 (FERM BP-7374), and *Pseudomonas jessenii* P161 (FERM BP-7376) to biosynthesize the polyhydroxyalkanoate from ω -(phenylmethyl)oxy]alkanoic acid expressed by chemical formula (19):



wherein x can be one or more integers within the range shown in the


chemical formula (19) as a raw material under a condition₁ which comprises the ω-[(phenylmethyl)oxy]alkanoic acid expressed by the chemical formula (19).

26. (Currently Amended) The method for producing a polyhydroxyalkanoate according to claim 25, wherein the polyhydroxyalkanoate comprises at least one unit expressed by ~~the following~~ chemical formulas (2) and (3):




wherein y and z can be one or more integers within the range shown in the chemical formulas (2) and (3), while being independent from the unit expressed by the chemical formula (1).

27. (Previously Presented) The method for producing a polyhydroxyalkanoate according to claim 25, wherein the ω-[(phenylmethyl)oxy]alkanoic acid expressed by said chemical formula (19) is 4-[(phenylmethyl)oxy]butyric acid expressed by chemical formula (23):




or 5-[(phenylmethyl)oxy]valeric acid expressed by chemical formula (24):



28. (Currently Amended) The method for producing a polyhydroxyalkanoate according to claim 25, comprising allowing the microorganism to biosynthesize ~~with an ability to produce~~ a polyhydroxyalkanoate comprising simultaneously, in at least a molecule thereof, the monomer unit of 3-hydroxy- ω -[(phenylmethyl)oxy]alkanoic acid expressed by chemical formula (1) and


a 3-hydroxy-alkanoic acid unit expressed by chemical formula (22):



wherein m can be one or more integers within the range shown in the chemical formula (22), and R₁₈ comprises a residue having either a phenyl structure or a thienyl structure, or

a 3-hydroxy- ω -cyclohexylalkanoic acid unit expressed by chemical formula


(5):



wherein R₁ is selected from the group consisting of H, CN, NO₂, halogen, CH₃, C₂H₅, C₃H₇, CF₃, C₂F₅ and C₃F₇, and k can be one or more integers within the range shown in the chemical formula (5),


from ω-[(phenylmethyl)oxy]alkanoic acid expressed by the chemical formula (19), ~~and~~ and

an alkanolic acid expressed by chemical formula (20):



wherein q can be one or more integers within the range shown in the chemical formula (20), and R₁₆ comprises a residue having either a phenyl structure or a thienyl structure, or


ω-cyclohexylalkanoic acid expressed by chemical formula (21):




wherein R₁₇ is selected from the group consisting of H, CN, NO₂, halogen, CH₃, C₂H₅, C₃H₇, CF₃, C₂F₅, and C₃F₇, and r can be one or more integers within the range shown in the chemical formula (21) as raw materials to biosynthesize the polyhydroxyalkanoate

under a condition, which comprises ω-[(phenylmethyl)oxy]alkanoic acid expressed by the chemical formula (19); and alkanoic acid expressed by the chemical formula (20) or ω-cyclohexylalkanoic acid expressed by the chemical formula (21),


wherein R₁₆ in chemical formula (20) and R₁₈ in chemical formula (22), i.e. residues having either a phenyl structure or a thienyl structure, are, independently, at least one group selected from the group consisting of residues




wherein R₁₉ is selected from the group consisting of H, halogen, CN, NO₂, CH₃, C₂H₅, C₃H₇, CH=CH₂, CF₃, C₂F₅, and C₃F₇, and in a case where there exist a plurality of units, R₁₉ may be different for each unit;




wherein R_4 is selected from the group consisting of H, halogen, CN, NO_2 , CH_3 , C_2H_5 , C_3H_7 , SCH_3 , CF_3 , C_2F_5 and C_3F_7 , and in a case where there exist a plurality of units, R_4 may be different for each unit;



wherein R_5 is selected from the group consisting of H, halogen, CN, NO_2 , CH_3 , C_2H_5 , C_3H_7 , CF_3 , C_2F_5 and C_3F_7 , and in a case where there exist a plurality of units, R_5 may be different for each unit;







wherein R_6 is selected from the group consisting of H, halogen, CN, NO_2 , CH_3 , C_2H_5 , C_3H_7 , $(\text{CH}_3)_2\text{-CH}$, $(\text{CH}_3)_3\text{-C}$, COOR_7 , and SO_2R_8 , ~~wherein~~ where R_7 represents any one selected from the group consisting of H, Na, K, CH_3 and C_2H_5 , and R_8 represents any one selected from the group consisting of OH, ONa, OK, halogen, OCH_3 and OC_2H_5 ; ~~CH_3 , C_2H_5 , C_3H_7 , $(\text{CH}_3)_2\text{-CH}$ and $(\text{CH}_3)_3\text{-C}$~~ , and in a case where there exist a plurality of units, R_6 may be different for each unit;




wherein R_9 is selected from the group consisting of H, halogen, CN, NO_2 , CH_3 , C_2H_5 , C_3H_7 , $(\text{CH}_3)_2\text{-CH}$, $(\text{CH}_3)_3\text{-C}$, COOR_{10} , and SO_2R_{11} , ~~wherein~~ where R_{10}

represents any one selected from the group consisting of H, Na, K, CH_3 , and C_2H_5 , and R_{11} represents any one selected from the group consisting of OH, ONa, OK, halogen, OCH_3 , and OC_2H_5 , CH_3 , C_2H_5 , C_3H_7 , $(\text{CH}_3)_2\text{-CH}$ and $(\text{CH}_3)_3\text{-C}$, and in a case where there exist a plurality of units, R_9 may be different for each unit;



wherein R_{12} is selected from the group consisting of H, halogen, CN, NO_2 , CH_3 , C_2H_5 , C_3H_7 , $(\text{CH}_3)_2\text{-CH}$, $(\text{CH}_3)_3\text{-C}$, COOR_{13} , and SO_2R_{14} (wherein where R_{13} represents any one selected from the group consisting of H, Na, K, CH_3 , and C_2H_5 , and R_{14} represents any one selected from the group consisting of OH, ONa, OK, halogen, OCH_3 ,

and OC_2H_5 , CH_3 , C_2H_5 , C_3H_7 , $(\text{CH}_3)_2\text{CH}$ and $(\text{CH}_3)_3\text{C}$, and in a case where there exist a plurality of units, R_{12} may be different for each unit; and



wherein R_{15} is selected from the group consisting of H, halogen, CN, NO_2 , CH_3 , C_2H_5 , C_3H_7 , $(\text{CH}_3)_2\text{CH}$, $(\text{CH}_3)_3\text{C}$, COOR_{16} , and SO_2R_{17} , ~~wherein~~ where R_{16} represents any one selected from the group consisting of H, Na, K, CH_3 , and C_2H_5 , and R_{17} represents any one selected from the group consisting of OH, ONa, OK, halogen, OCH_3 , and OC_2H_5 , CH_3 , C_2H_5 , C_3H_7 , $(\text{CH}_3)_2\text{CH}$ and $(\text{CH}_3)_3\text{C}$, and in a case where there exist a plurality of units, R_{15} may be different for each unit.

29. (Currently Amended) The method for producing a polyhydroxyalkanoate according to claim 25, wherein ~~said condition is that said microorganisms~~ the microorganism is cultured in a medium containing the ω -[(phenylmethyl)oxy]alkanoic acid expressed by the chemical formula (19).

30. (Currently Amended) The method for producing a polyhydroxyalkanoate according to claim 28, ~~wherein comprising culturing the microorganism~~ said condition is that said microorganism is cultured in a medium containing the ω -[(phenylmethyl)oxy]alkanoic acid expressed by the chemical formula (19)

and the alkanoic acid expressed by the chemical formula (20) or the ω -cyclohexylalkanoic acid expressed by the chemical formula (21).

31. (Currently Amended) The method for producing a polyhydroxyalkanoate according to claim 29, wherein said medium contains at least one selected from the group consisting of peptides, yeast extract, organic acids or salts thereof, amino acids or salts thereof, saccharides, and straight-chain alkanoic acids, which ~~is~~ are saturated or unsaturated fatty ~~acid~~ acids having 4 to 12 carbon atoms or salts thereof.

32. (Currently Amended) The method for producing a polyhydroxyalkanoate according to claim 31, wherein the peptide is polypeptone; the organic acids or salts thereof are one or more compounds selected from the group consisting of pyruvic acid, oxaloacetic acid, citric acid, isocitric acid, ketoglutaric acid, succinic acid, fumaric acid, malic acid, lactic acid, and salts thereof; the amino acids or salts thereof are one or more compounds selected from the group consisting of glutamic acid, aspartic acid, and salts thereof; and the saccharides are one or more compounds selected from the group consisting of glyceraldehyde, erythrose, arabinose, xylose, glucose, galactose, mannose, fructose, glycerol, erythritol, xylitol, gluconic acid, glucuronic acid and galacturonic acid, maltose, sucrose, and lactose.

33. (Currently Amended) The method for producing a

polyhydroxyalkanoate according to claim 29, wherein ~~said culture~~the culturing of the
~~microorganisms~~microorganism comprises two or more culturing steps.

34. (Currently Amended) The method for producing a
polyhydroxyalkanoate according to claim 33, wherein ~~said~~the culturing is performed
~~by culture~~ is a fed-batch culture.

35. (Currently Amended) The method for producing a
polyhydroxyalkanoate according to claim 29, comprising a step of recovering ~~a~~the
polyhydroxyalkanoate comprising the 3-hydroxy- ω -[(phenylmethyl)oxy]alkanoic acid unit
expressed by the chemical formula (1) generated by the microorganism from ~~the~~ cells of
the microorganism.

36-37. (Cancelled)